A commutative noetherian local ring of embedding dimension 4 having a transcendental series of Betti numbers

Jan-Erik Roos
Department of Mathematics
Stockholm University
e-mail: jeroos@math.su.se

February 14, 2017

arXiv:1702.04262v1 [math.AC] 14 Feb 2017

Abstract

We construct a ring with the properties of the title of the paper. We also construct some other local rings of embedding dimension 4 with exotic properties. Among the methods used are the Macaulay2-package DGAlgebras by Frank Moore, combined with and inspired by results by Anick, Avramov, Backelin, Katthän, Lemaire, Levin, Löfwall and others.

Mathematics Subject Classification (2010): Primary 13Dxx, 16Dxx, 68W30; Secondary 16S37, 55Txx

Keywords. local ring, Koszul complex, Yoneda Ext-algebra,
Hilbert series, Macaulay2, Betti series, Avramov spectral sequence

0. Introduction

The aim of the present paper is to prove the following (if V is a vector space over a field k we write $|V|$ instead of $\dim_k V$):

THEOREM 1 – *Let k be a field and $k[x,y,z,u]$ the polynomial ring in four variables of degree one and let*

$$R = \frac{k[x,y,z,u]}{(x^3, x^2y, (x+u)(y^2+z^2)+yzu, zu^2, u^3)} \tag{1}$$

be the graded quotient ring. Let

$$P_R(Z) = \sum_{i\geq 0} |\mathrm{Tor}_i^R(k,k)|Z^i$$

be the corresponding series of Betti numbers of R. Then $P_R(Z)$ is a transcendental function:

$$P_R(Z) = \frac{Z(1+Z)^2}{(1+Z)(1-Z-Z^2)^2\prod_{n=1}^{\infty}(1-Z^{2n}) - 1 + 2Z + 2Z^2 - 4Z^3 - 4Z^4}$$

If one replaces $k[x,y,z,u]$ with the formal power series ring $k[[x,y,z,u]]$ one obtains a local ring with the same homological properties. The Theorem might be of interest in itself (previously known examples of trancendental series needed ≥ 5 variables) but more interesting are probably the methods used to obtain the example (described in section 1 below) and the methods of proof described in section 2. There is now a possibility to obtain a new good insight in the theory of the homology of local rings of embedding dimension 4. In section 3 we will in particular indicate that the following ring which is a simpler variant of (1)

$$R = \frac{k[x,y,z,u]}{(x^3, x^2y, (x+u)(y^2+z^2), zu^2, u^3)} \tag{2}$$

has a $P_R(Z) = (1-Z^2)(1+Z)/(1-3Z-3Z^3-2Z^5)$ i.e. a rational function *but* is such that the Yoneda Ext-algebra $\mathrm{Ext}^*_R(k,k)$ is *not* finitely generated as an algebra. This last phenomenon was also previously only known in the embedding dimension ≥ 5 cases. We will also deduce some other transcendental results and also show that there are quotients of $k[x,y,z,u]$ with an ideal with only *six* cubic generators which is non-Golod but has the multiplication in Koszul homology equal to zero (the first examples of this last phenomenon with more relations - monomials of higher degrees and more variables - were obtained by Lukas Katthän [KAT]).

1. How the example in Theorem 1 was found

In [A] (announced in 1980 [A-CR]), David Anick published the first example (example 7.1 page 29 of [A]) of an R in 5 variables and 7 quadratic relations for which $P_R(Z)$ was transcendental. Here is a variant of his example with only 5 quadratic relations which has the same property and which will be useful for us:

$$R = \frac{k[x_1,x_2,x_3,x_4,x_5]}{(x_1^2, x_1x_2, x_1x_3+x_2x_4+x_3x_5, x_4x_5, x_5^2)} \tag{3}$$

In this case the Hilbert series of R is $H(t) = (1+3t+t^2-2t^3+t^4)/(1-t)^2$ and the Hilbert series of the Koszul dual $R^!$ of R is

$$R^!(t) = \frac{(1+t)^2}{(1-t-t^2)^2} \prod_{n=1}^{\infty} \frac{1+t^{2n-1}}{1-t^{2n}}$$

and

$$1/P_R(Z) = (1+1/Z)/R^!(Z) - H(-Z)/Z \tag{4}$$

Remark: For the case studied by Anick [A] we had that $H(t) = 1 + 5t + 7t^2$ and

$$R^!(t) = \frac{1}{(1-2t)^2} \prod_{n=1}^{\infty} \frac{1+t^{2n-1}}{1-t^{2n}}$$

and the formula (4) was also valid in his case.

We now address the problem of constructing a 4-variable version of the ring (3). We start with the ring $S_2 = k[x,y,z,u]/(x^2, xy, zu, u^2)$ where x, y, z, u correspond to x_1, x_2, x_4, x_5 and try to add an extra relation corresponding to $x_1x_3 + x_2x_4 + x_3x_5$. How to do this is not at all evident and instead we try to add a new relation which is a linear combination with coefficients 0 or 1 of the 6 nonzero quadratic monomials yu, xu, z^2, yz, xz, y^2 in S_2. There are 64 such linear combinations but each of them leads to a rational $P_R(Z)$ (they all occur in the Appendix to [R4]). Instead we turn to the study of cubic relations, i.e. we start with a new ring

$$S = \frac{k[x,y,z,u]}{(x^3, x^2y, zu^2, u^3)}$$

and try to add a linear combination (coefficients 0 or 1) of the 16 non-zero cubic monomials

$$yu^2, xu^2, z^2u, yzu, xzu, y^2u, xyu, x^2u, z^3, yz^2, xz^2, y^2z, xyz, x^2z, y^3, xy^2$$

in S. There are $2^{16} = 65536$ cases to study and this can be done automatically using the following input programme to Macaulay 2 which was written at my request several years ago by Mike Stillman (it is an elegant version of a programme I wrote for Macaulay 1 a long time ago (cf. [R3, pp. 294-296])

```
binaries = (n) -> (
if n === 0 then {}
else if n === 1 then {{0},{1}}
else (
r := binaries(n-1);
join(apply(r, i->prepend(0,i)),
apply(r, i->prepend(1,i)))))
doit = (i) -> (
h := hypers_(0,i);
J1 := J + ideal(h);
<< newline << flush;
<< ``--- n = `` << i << `` ideal = `` << hypers_(0,i) << `` ---'' << flush;
<< newline << flush;
```

```
E := res J1;
<< newline << flush;
<< `` `` << betti E << flush;
<< newline << flush;
A := (ring J1)/J1;
C := res(coker vars A, LengthLimit=>6);
<< newline << flush;
<< `` `` << betti C << flush;
<< newline << flush;
<< `` `` <<hilbertSeries(A,Order=>12)<< flush;
<< newline << flush;
)
makeHyperplanes = (J) -> (
I := matrix basis(3,coker gens J);
c := numgens source I;
m := transpose matrix binaries c;
I * m)
R = QQ[x,y,z,u]
J = ideal(x^3,x^2*y,z*u^2,u^3)
time hypers = makeHyperplanes J;
time scan(numgens source hypers, i -> doit i);
```

We obtain 20 possibilities for the $P_R(z)$ up to degree 6 presented here in increasing order (of course there can be variations inside each case $P_i(z)$ due e.g. to different Koszul homology) :

$P_1(z)$=$1+4z+11z^2+27z^3+62z^4+137z^5+295z^6+\ldots$ $P_2(z)$=$1+4z+11z^2+27z^3+64z^4+152z^5+364z^6+\ldots$

$P_3(z)$=$1+4z+11z^2+28z^3+69z^4+168z^5+407z^6+\ldots$ $P_4(z)$=$1+4z+11z^2+28z^3+69z^4+169z^5+414z^6+\ldots$

$P_5(z)$=$1+4z+11z^2+28z^3+69z^4+172z^5+431z^6+\ldots$ $P_6(z)$=$1+4z+11z^2+28z^3+70z^4+173z^5+427z^6+\ldots$

$P_7(z)$=$1+4z+11z^2+28z^3+70z^4+177z^5+451z^6+\ldots$ $P_8(z)$=$1+4z+11z^2+29z^3+74z^4+188z^5+476z^6+\ldots$

$P_9(z)$=$1+4z+11z^2+29z^3+75z^4+193z^5+496z^6+\ldots$ $P_{10}(z)$=$1+4z+11z^2+29z^3+75z^4+193z^5+498z^6+\ldots$

$P_{11}(z)$=$1+4z+11z^2+29z^3+75z^4+194z^5+503z^6+\ldots$ $P_{12}(z)$=$1+4z+11z^2+30z^3+80z^4+213z^5+567z^6+\ldots$

$P_{13}(z)$=$1+4z+11z^2+30z^3+81z^4+218z^5+587z^6+\ldots$ $P_{14}(z)$=$1+4z+11z^2+30z^3+82z^4+224z^5+612z^6+\ldots$

$P_{15}(z)$=1+4z+11z^2+31z^3+86z^4+238z^5+660z^6+... $P_{16}(z)$=1+4z+11z^2+31z^3+87z^4+244z^5+685z^6+...

$P_{17}(z)$=1+4z+11z^2+31z^3+88z^4+249z^5+705z^6+... $P_{18}(z)$=1+4z+11z^2+32z^3+92z^4+263z^5+755z^6+...

$P_{19}(z)$=1+4z+11z^2+32z^3+93z^4+269z^5+780z^6+... $P_{20}(z)$=1+4z+11z^2+33z^3+99z^4+294z^5+877z^6+...

Which one of these cases could give a transcendental $P_R(Z)$?

One can guess a possibility by studying how the transcendental $P_R(z)$ comes up in five variables when we add linear combinations of quadratic monomials to the ring in 5 variables: $\mathbf{Q}[x,y,z,u,v]/(x^2,xy,uv,v^2)$ i.e. modifying the previous input programme by replacing R by $QQ[x,y,z,u,v]$, J by $(x^2, x*y, u*v, v^2)$ and the line

```
I := matrix basis(3,coker gens J);
```

by the line

```
I := matrix basis(2,coker gens J);
```

There are now "only" $2^{11} = 2048$ cases to study and now we obtain only 10 possibilities for the $P_R(z)$ up to degree 6 presented here in increasing order:

$P_1(z)$=1+5z+15z^2+37z^3+82z^4+170z^5+337z^6+... $P_2(z)$=1+5z+15z^2+38z^3+88*z^4+192z^5+406z^6+...

$P_3(z)$=1+5z+15z^2+38z^3+89z^4+199z^5+432z^6+... $P_4(z)$=1+5z+15z^2+38z^3+91z^4+216z^5+516z^6+...

$P_5(z)$=1+5z+15z^2+39z^3+95z^4+222z^5+506z^6+... $P_6(z)$=1+5z+15z^2+39z^3+96z^4+231z^5+553z^6+...

$P_7(z)$=1+5z+15z^2+39z^3+97z^4+237z^5+575z^6+... $P_8(z)$=1+5z+15z^2+39z^3+99z^4+254z^5+659z^6+...

$P_9(z)$=1+5z+15z^2+40z^3+104z^4+268z^5+689z^6+... $P_{20}(z)$=1+5z+15z^2+41z^3+112z^4+306z^5+836z^6+...

There are e.g. 1024 cases of an extra quadratic relation corresponding to $P_1(z)$ and 576 cases corresponding to $P_4(z)$, but only 8 for $P_2(z)$ including the case of the extra relation $xz+yu+zv$ leading to the transcendental series mentioned in the introduction (the other 7 cases of $P_2(z)$ lead to the same total series). Furthermore, there are also 8 cases for $P_6(z)$ and they all lead to cases where the Yoneda Ext-algebra $\mathrm{Ext}_R^*(k,k)$ is not finitely generated but the series of Betti numbers is rational (this phenomenon was first found by me in [R1] (inspired by Lemaire [LEM]).

We now try to use this information to try to guess what happens for the 65536 embedding dimension 4 cases above. In this situation there are only 80 cases of $P_{10}(z)$ and only two of them correspond to adding a cubic relation with is a sum of 5 cubic monomials (the other cases need more monomials): case 9257 which corresponds to $xy^2+xz^2+y^2u+yzu+z^2u$ and case 13345 which corresponds to $xy^2+xyz+xz^2+y^2u+z^2u$. Here one of

these cases can be obtained from the other one by permuting x and u. We can therefore restrict ourselves to the homological study of the ring

$$\frac{k[x,y,z,u]}{(x^3, x^2y, (x+u)(y^2+z^2)+yzu, zu^2, u^3)} \tag{1}$$

mentioned in the introduction and which will be the subject of the next section.

2. The treatment of the example in Theorem 1

We will now use the programme `DGAlgebra` of [MOO] on the ring (1).

The Koszul homology of the ring has the Betti numbers given by Macaulay:

total:	1	5	12	12	4
0:	1	.	.	.	.
1:	.	.	.	.	.
2:	.	5	2	.	.
3:	.	.	1	.	.
4:	.	.	8	10	3
5:	.	.	.	.	.
6:	.	.	1	2	1

From this we can guess that the multiplication in $HKR = \mathrm{Tor}_*^{k[x,y,z,u]}(k[x,y,z,u]/J, k)$, where J is the ideal in (1) is rather complicated. We determine details about that multiplication as a preparation for using the Avramov spectral sequence ([AV0],[AV1]) where $\tilde{R} = k[x,y,z,u]$ and $R = \tilde{R}/J$:

$$E^2_{p,q} = \mathrm{Tor}^{HKR}_{p,q}(k,k) => \mathrm{Tor}^R_*(k,k)//\mathrm{Tor}^{\tilde{R}}_*(k,k) \tag{5}$$

by using the following infile for Macaulay:

```
loadPackage(``DGAlgebras'')
R=QQ[x,y,z,u]/ideal(x^3,x^2*y,z*u^2,u^3,x*y^2+x*z^2+y^2*u+y*z*u+z^2*u)
res(ideal R); betti oo
res(coker vars R,LengthLimit => 6); betti oo
HKR=HH koszulComplexDGA(R)
generators HKR
for n from 1 to length(generators HKR) list degree X_n
ideal HKR
```

```
I=ideal(vars HKR)
I^2; trim(oo)
I^3; trim(oo)
res(coker vars HKR,LengthLimit => 3);
betti oo
res(coker vars HKR,LengthLimit => 4);
betti oo
```

From the Macaulay output of this input file we see that the augmentation ideal I of HKR satisfies $I^3 = 0$, and that I^2 is generated by

$$X_4X_5, X_2X_5, X_1X_5, X_3X_4, X_2X_4, X_1X_4, X_2X_3, X_4X_7, X_2X_7, X_1X_7, X_5X_6, X_4X_6, X_3X_6. X_6X_7$$

Therefore we will be able to calculate the left side of the Avramov spectral sequence (5) exactly as in [R4], where we used the Theorem 1.3 on page 310 of Clas Löfwall's paper [L1], provided we can determine the Hilbert series $HKR^!(z,z)$ of the Koszul dual $HKR^!$ of HKR and the Hilbert series $HKR(-z,z)$ of HKR. From the output file we see that HKR can be presented as follows: It has five generators X_1, X_2, X_3, X_4, X_5 of degree (1,3),two generators X_6, X_7 of degree (2,4), three generators X_8, X_9, X_{10} of degrees (2,5),(2,6),(2,8) respectively, four generators $X_{11}, X_{12}, X_{13}, X_{14}$ of degree (3,7), two generators X_{15}, X_{16} of degree (3,9), two generators X_{17}, X_{18} of degree (4,8) and one generator X_{19} of degree (4,10).

Furthermore the first five variables X_1, X_2, X_3, X_4, X_5 skew-commute, the variables $X_6, X_7, X_8, X_9, X_{10}$ commute among themselves and with X_1, X_2, X_3, X_4, X_5, etc.

The relations are as follows: The product of the last 12 variables $X_8, \dots X_{19}$ with all the 19 variables $X_1 \dots X_{19}$ are 0. For the first five generators X_1, X_2, X_3, X_4, X_5 we only have the following three quadratic relations: $X_3X_5, X_1X_3 + X_2X_4 - X_4X_5, X_1X_2$, and if we also take into account the additional variables X_6, X_7 we have the extra quadratic relations $X_1X_6, X_2X_6, X_3X_7, X_5X_7, X_6X_6, X_7X_7$. It follows that the if

(6)

$$C = \frac{k[X_1, X_2, X_3, X_4, X_5, X_6, X_7]}{(X_3X_5, X_1X_3 + X_2X_4 - X_4X_5, X_1X_2, X_1X_6, X_2X_6, X_3X_7, X_5X_7, X_6X_6, X_7X_7)}$$

then the Koszul dual $HKR^!$ of HKR can be presented as a coproduct

$$C^! \sqcup k < x_8, x_9, x_{10}, x_{11}, x_{12}, x_{13}, x_{14}, x_{15}, x_{16}, x_{17}, x_{18}, x_{19} > \tag{7}$$

where the second algebra in the coproduct is the free algebra on the dual generators $x_8, \dots x_{19}$ of $X_8, \dots X_{19}$.

Furthermore, it is well-known (cf. e.g. [LEM]) that for the coproduct $A \sqcup B$ of two graded connected algebras, we have the following formula for the respective Hilbert series

$$\frac{1}{(A \sqcup B)(z)} + 1 = \frac{1}{A(z)} + \frac{1}{B(z)}$$

(this works also for the bigraded case) and it follows that

$$\frac{1}{HKR^!(z,z)} + 1 = \frac{1}{C^!(z,z)} + 1 - 3z^3 - 6z^4 - 3z^5 \tag{8}$$

and we have therefore reduced our problem to calculate $C^!(z,z)$ where C only involves the first 7 variables $X_1, X_2, X_3, X_4, X_5, X_6, X_7$

We start with an algebra D that involves only the first 5 variables X_1, X_2, X_3, X_4, X_5:

$$D = \frac{k[X_1, X_2, X_3, X_4, X_5]}{(X_3X_5, X_1X_3 + X_2X_4 - X_4X_5, X_1X_2)}$$

The variables skewcommute. The Koszul dual $D^!$ is the quotient of the free algebra in the dual variables x_i of the X_i :

$$D^! = \frac{k < x_1, x_2, x_3, x_4, x_5 >}{([x_1, x_4], [x_1, x_5], [x_2, x_3], [x_2, x_5], [x_3, x_4], [x_1, x_3] + [x_4, x_5], [x_2, x_4] + [x_4, x_5]}$$

where the $[x_i, x_j]$ are Lie commutators. From this one sees exactly as in Löfwall's and mine study of the Anick example [LR1] that the underlying Lie algebra g_D of $D^!$ sits in the middle of an extension

$$0 \longrightarrow \oplus_{i=1}^{\infty} a_i \longrightarrow g_D \longrightarrow f_1 \times f_2 \longrightarrow 0$$

i.e. is the extension of the product of two free Lie algebras f_1 generated by x_1 and x_2 and f_2 generated by x_3 and x_5 with the infinite abelian Lie algebra

$$a = \oplus_{i=1}^{\infty} a_i$$

where the a_i are onedimensional generated by $a_1 = x_4$, $a_2 = [x_5, x_4]$, $a_3 = [x_5, [x_5, x_4]]$ etc. which commute and by the 2-cocycle $\gamma : f_1 \times f_2 \to a$ defined by $\gamma(x_1, x_3) = -\gamma(x_3, x_1) = a_2$ and $\gamma(x_2, x_5) = -\gamma(x_5, x_2) = a_2$. It follows that

$$D^!(z) = \frac{1}{(1-2z)^2} \prod_{n=1}^{\infty} \frac{1}{(1-z^n)} \tag{7}$$

But since we need $D^!(z,z)$ we have to replace the z by z^2 in the formula (9) to get $D^!(z,z)$. We next want to incorporate the variables X_6, X_7 (which *commute* with the X_1, X_2, X_3, X_4, X_5) into the picture. It turns out that we have to study the underlying Lie algebra of the Koszul dual $C^!$ of the C in (6), i.e. the quotient of

$$k < x_1, x_2, x_3, x_4, x_5, x_6, x_7 >$$

with the ideal generated by

$$[x_1, x_4], [x_1, x_5], [x_2, x_3], [x_2, x_5], [x_3, x_4], [x_1, x_3] + [x_4, x_5], [x_2, x_4] + [x_4, x_5],$$

as before and the extra ideal generators:

$$[x_1, x_7], [x_2, x_7], [x_4, x_7], [x_6, x_7], [x_3, x_6], [x_4, x_6], [x_6, x_7]$$

where x_6 and x_7 are dual to X_6 and X_7. From the preceding presentation one sees that the Lie algebra g_C of $C^!$ sits in the middle of an extension:

$$0 \longrightarrow \oplus_{i=1}^{\infty} a_i \longrightarrow g_C \longrightarrow F_1 \times F_2 \longrightarrow 0 \tag{10}$$

where F_1 is the free Lie algebra generated by x_1, x_2, x_6 of degrees $1, 1, 2$ and F_2 is the free Lie algebra generated by x_3, x_5, x_7 of degrees $1, 1, 2$ and the a_i:s are as before, and the x_6 and x_7 operate in the trivial way on the a_i. Furthermore we have a new cocycle

$$\Gamma : F_1 \times F_2 \longrightarrow \oplus_{i=1}^{\infty} a_i$$

where as before Γ is defined by its value on the generators by $-\Gamma(x_3, x_1) = \Gamma(x_1, x_3) = a_2$ and $-\Gamma(x_5, x_2) = \Gamma(x_2, x_5) = a_2$ and being zero for all other pairs of generators. By calculating $H^2(F_1 \times F_2, a)$ for (8) one sees that the relations are those of $C^!$ and therefore the Hilbert series

$$C^!(z) = \frac{1}{(1-2z-z^2)^2} \prod_{n=1}^{\infty} \frac{1}{(1-z^n)} \tag{11}$$

if the variables $x_1, x_2, x_3, x_4, x_5, x_6, x_7$ are given the degrees $1, 1, 1, 1, 1, 2, 2$. But for the calculation of the series of $HKR^!(z,z)$ these variables should be given the degrees $2, 2, 2, 2, 2, 3, 3$ so that we get the series

$$\frac{1}{(1-2z^2-z^3)^2} \prod_{n=1}^{\infty} \frac{1}{(1-z^{2n})} \tag{12}$$

Using the formulae (8) and (12) we obtain

$$\frac{1}{HKR^!(z,z)} = (1-2z^2-z^3)^2\prod_{n=1}^{\infty}(1-z^{2n}) - 3z^3 - 6z^4 - 3z^5 \tag{13}$$

But $I^3 = 0$ and

$$I^2=(X_4X_5, X_2X_5, X_1X_5, X_3X_4, X_2X_4, X_1X_4, X_2X_3, X_4X_7, X_2X_7, X_1X_7, X_5X_6, X_4X_6, X_3X_6. X_6X_7)$$

and this gives that $HKR(x,y) = 1 + 5xy + 5xy^2 + 6xy^3 + 3xy^4 + 7x^2y^2 + 6x^2y^3 + x^2y^4$, so that

$$HKR(-z,z) = 1 - 5z^2 - 5z^3 + z^4 + 3z^5 + z^6 \tag{14}$$

and finally the formula

$$\frac{1}{P_{HKR}(z,z)} = (1+1/z)/HKR^!(z,z) - HKR(-z,z)/z$$

gives that

$$\frac{1}{P_{HKR}(z,z)} = (1+1/z)[(1-2z^2-z^3)^2\prod_{n=1}^{\infty}(1-z^{2n}) - 3z^3 - 6z^4 - 3z^5)]$$

$$-(1-5z^2-5z^3+z^4++3z^5+z^6)/z$$

and *if the Avramov spectral sequence degenerated* this series should be the same as $(1+z)^4/P_R(z)$ so the final result should be

$$P_R(z) = \frac{z(1+z)^2}{(1+z)(1-z-z^2)^2\prod_{n=1}^{\infty}(1-z^{2n}) - 1 + 2z + 2z^2 - 4z^3 - 4z^4}$$

$$= 1 + 4z + 11z^2 + 29z^3 + 75z^4 + 193z^5 + 498z^6 + 1289z^7 + 3341z^8 + 8663z^9 + 22466z^{10} + \dots$$

as asserted in our Theorem 1. The following calculation in Macaulay2

```
res(coker vars R, LengthLimit => 11); betti oo
```

gives the diagram of Betti numbers (16) below whose first row (the total Betti numbers) gives support to this assertion. But the Theorem 5.9 of [AV0] gives first that the differentials d^r of the Avramov spectral sequence are 0 for $r > 2$ but furthermore since the dimensions of the two sides of the Avramov spectral sequence are the same in degrees ≤ 10

it follows from the proof of Theorem 5.9 in [AV0] and the structure of the matrix Massey products that there are no non-zero differentials, and Theorem 1 is proved. We can also get a bigraded more precise version of all this: Indeed, we also have a series

$$P_R(x,y) = \sum_{p,q} |\mathrm{Tor}^R_{p,q}(k,k)| x^p y^q$$

which takes into account the extra grading of R (we have that $P_R(z,1)$ is the old $P_R(z)$) given as

$$P_R(x,y) = \frac{(1+xy)^4}{(1+1/x)/A - H/x} \tag{15}$$

where

$$1/A = (1-2x^2y^3-x^3y^4)^2 \prod_{n=1}^{\infty}(1-x^{2n}y^{3n}) - x^3y^5 - x^3y^6 - x^3y^8 - 4x^4y^7 - 2x^4y^9 - 2x^5y^8 - x^5y^{10}$$

and H is equal to

$$1 - 5x^2y^3 - 2x^3y^4 - x^3y^5 - x^3y^6 - x^3y^8 - 4x^4y^7 - 2x^4y^9 - 2x^5y^8 - x^5y^{10} + 7x^4y^6 + 6x^5y^7 + x^6y^8$$

and the expansion of $P_R(x,y)$ of (15) up to degree 11 gives the same diagram of graded Betti numbers that was earlier found by the Macaulay2 calculation:

(16)

total:	1	4	11	29	75	193	498	1289	3341	8663	22466	58264
0:	1	4	6	4	1	.	.	.	.	.	.	.
1:	.	.	5	22	38	32	13	2	.	.	.	.
2:	.	.	.	1	22	92	171	169	92	26	3	.
3:	.	.	.	1	8	34	135	389	689	744	493	196
4:	.	.	.	.	.	10	87	346	923	1870	2835	3066
5:	.	.	.	1	6	15	22	96	645	2426	5739	9705
6:	.	.	.	.	.	10	65	186	348	964	4302	14387
7:	.	.	.	.	.	.	2	73	459	1446	3296	8595
8:	.	.	.	.	.	.	2	20	112	669	3025	9538
9:	.	.	.	.	.	.	.	.	30	320	1668	6469
10:	.	.	.	.	.	.	1	8	28	62	396	3282
11:	.	.	.	.	.	.	.	.	15	129	510	1377
12:	.	.	.	.	.	.	.	.	.	3	153	1278
13:	.	.	.	.	.	.	.	.	.	3	36	246
14:	.	.	.	.	.	.	.	.	.	.	.	60
15:	.	.	.	.	.	.	.	.	.	1	10	45
16:	.	.	.	.	.	.	.	.	.	.	.	20

3. Other embedding dimension 4 cases

The example we have just studied is certainly not unique. Let us illustrate this with two examples: If one adds the relation z^3 to the example in Theorem 1 we obtain the following

$$R_a = \frac{k[x,y,z,u]}{(x^3, x^2y, (x+u)(y^2+z^2)+yzu, z^3, zu^2, u^3)}$$

which has Hilbert series

$$\frac{1+3T+6T^2+4T^3-T^4-7T^5}{1-T}$$

Furthermore the homology of the Koszul complex of R_a is

total:	1	6	18	20	7
0:	1	.	.	.	.
1:	.	.	.	.	.
2:	.	6	2	.	.
3:	.	.	2	.	.
4:	.	.	14	20	7

We get same Hilbert series and the same diagram of Koszul homology if we replace z^3 with y^3+z^3 to get the ring

$$R_b = \frac{k[x,y,z,u]}{(x^3, x^2y, (x+u)(y^2+z^2)+yzu, y^3+z^3, zu^2, u^3)}$$

Both these ring have transcendental series of Betti numbers. These two series can be determined and they are different. For the case of R_a we have using `DGAlgebras` that HKR_a is generated by 6 skew-commuting variables $X_1, X_2, X_3, X_4, X_5, X_6$ of degree (1,3) (and 27 variables of higher degrees). The first 6 variables satisfy the following relations $X_5X_6, X_3X_6, X_3X_5, X_1X_3+X_2X_4-X_4X_6, X_1X_2$ and the Hilbert series of the corresponding Koszul dual is

$$\frac{1}{(1-2z)(1-3z)\prod_{n=1}^{\infty}(1-z^n)}$$

On the other hand, using `DGAlgebras` for R_b we find that HKR_b is still generated by 6 variables $X_1, X_2, X_3, X_4, X_5, X_6$ of degree (1,3) (but only 24 variables of higher degrees). In this case the first 6 variables satisfy the following relations $X_3X_6, X_1X_3+X_2X_4-X_4X_6, X_1X_2$ and in this case the Hilbert series of the corresponding Koszul dual is

$$\frac{1}{(1-2z)^2(1-z)\prod_{n=1}^{\infty}(1-z^n)}$$

Now we can continue our analysis as in section 2 and we obtain the following formulae:

$$P_{R_a}(Z) = \frac{Z(1+Z)^2}{(1-Z-Z^2)(1-3Z^2-Z^3)\prod_{n=1}^{\infty}(1-Z^{2n}) - 1 + 2Z + 3Z^2 - 6Z^3 - 7Z^4}$$

and

$$P_{R_b}(Z) = \frac{Z(1+Z)^2}{(1-Z)(1-2Z^2-Z^3)^2\prod_{n=1}^{\infty}(1-Z^{2n}) - 1 + 2Z + 3Z^2 - 6Z^3 - 7Z^4}$$

Note also that we could have started with many more examples in embedding dimension 4 than just our variation of the Anick case that we took in section 1.

Now let us also briefly analyze the case when

$$S = \frac{k[x,y,z,u]}{(x^3, x^2y, (x+u)(y^2+z^2), zu^2, u^3)}$$

Using `DGAlgebras` we obtain as before that

$$P_S(Z) = \frac{(1-Z)(1+z)^2}{1-3Z+3Z^3-2Z^5}$$

but we can also prove that neither $\mathrm{Ext}^*_S(k,k)$ nor $\mathrm{Ext}^*_{HKS}(k,k)$ are finitely generated as algebras equipped with the Yoneda product. The case S is the 4-variable of the 5-variable $k[x,y,z,u,v]/(x^2, xy, xz+zv, uv, v^2)$, a variant of which we treated in [R1], inspired by Lemaire [LEM]. An alternative way to treat S is to observe that the quotient map

$$\frac{k[x,y,z,u]}{(x^3, x^2y, zu^2, u^3)} \longrightarrow S$$

is a Golod map [LEV] so that $k[x,y,z,u] \to S$ is a composite of three Golod maps, and use [R2].

All this leads support to the surmise that the embedding dimension 4 case could be equally complicated as the general case, and one could even pose the

QUESTION 1: Let $\mathcal{E}4$ be the set of series $\sum_{n\geq 0} |\mathrm{Tor}^R_n(k,k)|Z^n$ for (R,m) a local commutative noetherian ring of embedding dimension ≤ 4 where $k = R/m$. Can $\mathcal{E}4$ be added to the list of 17 series of [A-Gu] that are all rationally related ?

One would certainly get a smaller set than $\mathcal{E}4$ if one restricted oneself to rings of the form $k[x,y,z,u]/(f_1, \ldots f_s)$ where the relations f_i were of degree ≤ 3 or ≤ 4 .

So far no embedding dimension 4 variant of the rings in [LR2] has been found. But even in this restricted case of degree 3 relations one can get surprizes: The ring

$$S_I = Q[x,y,z,u]/(u^3, xy^2, (x+y)z^2, x^2u + zu^2, y^*u + xzu, y^2z + yz^2)$$

constructed in [R5] (inspired by Lukas Katthän [KAT]) which has only six relations of degree 3, has as homology of the Koszul complex:

Total:	1	6	12	9	2
0:	1	.	.	.	.
1:	.	.	.	.	.
2:	.	6	.	.	.
3:	.	.	12	5	1
4:	.	.	.	4	.
5:	.	.	.	.	1

Furthermore the multiplication of elements of positive degree in this homology is zero (use `DGAlgebra`), and

$$\frac{(1+z)^4}{(1-6z^2-12z^3-9z^4-2z^5)} = 1+4z+12z^2+40z^3+130z^4+422z^5+1376z^6+4476z^7+\dots$$

whereas the series of the total Betti numbers of S_I is given by Macaulay2 as:

$$1+4z+12z^2+40z^3+130z^4+421z^5+1371z^6+4454z^7+\dots$$

so that S_I is *not* a Golod ring. In [R5] we show that if one accepts relations of degree 3 *and* 4, there are hundreds of such exotic non-Golod rings.

QUESTION 2: For local rings R of embedding dimension 4 with only cubic relations, are my R in Theorem 1 and the R_a and R_b essentially the only examples where we have a transcendental $P_R(Z)$?

4. References

[**A**] D.J. Anick, *A counterexample to a conjecture of Serre*, Ann. of Math. **115**, 1982, pp. 1-33; *Correction*, Ann. of Math. **116** , p. 661.

[**A-CR**] D.J. Anick, *Construction d'espaces de lacets et d'anneaux locaux à séries de Poincar-Betti non rationnelles*, C.R. Acad. Sc. Paris **290**, (1980), pp. A733-A736.

[**A-Gu**] D.J. Anick and Tor H. Gulliksen, *Rational dependence among Hilbert and Poincare series*, Journal of Pure and Applied Algebra **38**, 1985, pp. 135-157.

[**AV0**] Avramov, L. L., *The Hopf algebra of a local ring. (Russian)*, Izv. Akad. Nauk SSSR Ser. Mat. **38**, (1974), pp. 253-277. English translation [Math. USSR-Izv.**8** (1974), 259-284].

[**AV1**] Avramov, L. L., *Obstructions to the existence of multiplicative structures on minimal free resolutions.* Amer. J. Math., **103** (1981), pp. 1-31.

[**KAT**] L. Katthän, *A non-Golod ring with a trivial product on its Koszul homology*, `http://arxiv.org/pdf/1511.04883.pdf`

[**LEM**] Lemaire, Jean-Michel, *Algèbres connexes et homologie des espaces de lacets*, Lecture Notes in Mathematics, **422**, Springer-Verlag, Berlin-New York, 1974.

[**LEV**] Levin, Gerson, *Local rings and Golod homomorphisms*, Journal of Algebra **37**, 1975, pp 266289.

[**L1**] Löfwall, C., *On the subalgebra generated by one-dimensional elements in the Yoneda* Ext*–algebra*, in Algebra, algebraic topology, and their interactions, (J.–E. Roos, ed), Lecture Notes in Math., vol.**1183**, Springer-Verlag, Berlin–New York, 1986, pp. 291-338.

[**LR1**] C. Löfwall and J.-E. Roos, *Cohomologie des algèbres de Lie graduées et séries de Poincaré-Betti non rationnelles*, C.R. Acad. Sc. Paris **290**, 1980, pp. A733-A736.

[**LR2**] C. Löfwall and J.-E. Roos, *A nonnilpotent 1-2-presented graded Hopf algebra whose Hilbert series converges in the unit circle*, Adv. Math.**130**, 1997, pp. 161-200.

[**MOO**] Moore, Frank, `DGAlgebras.` A package for `Macaulay2` `http://www.math.uiuc.edu/Macaulay2/Packages/`

[**R1**] J.-E. Roos, *Relations between the Poincaré-Betti series of Loop Spaces and of Local rings*, Springer Lecture Notes in Math.**740**,1979, 285-322.

[**R2**] J.-E. Roos, *On the use of graded Lie algebras in the theory of local rings*, Commutative algebra: Durham 1981 (R. Y. S harp, ed.) London Math. Soc. Lecture Notes Ser. vol. 72, Cambridge Univ. Press, Cambridge, 1982, pp. 204230.

[**R3**] J.-E. Roos, *A computer-aided study of the graded Lie-algebra of a local commutative noetherian ring* (with an Appendix by Clas Löfwall), Journal of Pure and Applied Algebra **91** , 1994, pp. 255-315.

[**R4**] J.-E. Roos, *Homological properties of the homology algebra of the Koszul complex of a local ring: Examples and questions*, Journal of Algebra **465**, 2016, pp. 399-436.

[**R5**] J.-E. Roos, *On some unexpected rings that are close to Golod rings*, in preparation.